

The effects of reduced salinity on lobster (*Homarus americanus* Milne-Edwards) metabolism: implications for estuarine populations

Steven H. Jury*, Michael T. Kinnison, W. Hunting Howell,
Winsor H. Watson, III**

Zoology Department, Center for Marine Biology, University of New Hampshire, Durham, NH 03824, USA

(Received 3 March 1993; revision received 8 September 1993; accepted 15 October 1993)

Abstract

During periods of substantial freshwater runoff, lobsters that inhabit estuaries, such as the Great Bay Estuary in NH, are exposed for several days to weeks to seawater that is diluted as low as 10 ppt. To assess the physiological stress imposed by these conditions, we measured the oxygen consumption, heart rate, ventilation rate and hemolymph osmolarity of lobsters while sequentially exposing them, for 24-h periods, to seawater of 20, 15, and 10 ppt. Measurements of hemolymph osmolarity confirmed previous results which demonstrated that at salinities below 20 ppt lobsters are limited osmoregulators; allowing their hemolymph osmolarity to drop as the environmental salinity is reduced, but always maintaining it higher than the ambient osmolarity. All animals exposed to 10 ppt, at 15 °C, were capable of surviving for at least 72 h. There was a nearly linear increase in oxygen consumption, heart and scaphognathite rates in animals exposed to dilute seawater, with almost a twofold increase in metabolic rate when animals were moved from 20 to 15 to 10 ppt. At the lowest salinity tested (10 ppt) the average oxygen consumption was higher for females than for males. We conclude that at low salinities the energetic demands of osmoregulation are greater for females than males, and for both sexes the physiological stress imposed may determine, in part, their distribution and/or movements in estuarine habitats.

Key words: Estuary; Lobster; Metabolism; Oxygen consumption; Respirometer; Salinity

* Current address: N-ORCA, SSMC4, NOAA, Ninth Floor, 1305 East West Highway, Silver Spring, MD 20910, USA.

** Corresponding author.

1. Introduction

American lobsters, *Homarus americanus* Milne-Edwards are considered to be poor osmoregulators (Dall, 1980). Nonetheless, their range extends into many bays and estuaries along the North Atlantic coast (Thomas & White, 1969; Briggs & Mushacke, 1979; Munro & Therriault, 1983; Reynolds & Casterlin, 1985; Estrella & McKiernan, 1989; Vetrovs, 1990; Howell & Watson, 1991; Maynard, 1991; Robichaud & Campbell, 1991; Watson & Howell, 1991). These habitats are characterized by large seasonal fluctuations in salinity due to heavy rains and runoff from melting snow, which limits occupation of these areas to species that have evolved appropriate behavioral and/or physiological adaptations (Sanders et al., 1965; Theede, 1975; Vernberg & Vernberg, 1975; Lassere, 1976).

In the upper Great Bay Estuary, N.H., the salinity drops to between 10 and 15 ppt each spring (Nelson et al., 1981, 1982; Loder et al., 1983), which is close to the lethal limit for adult lobsters (McLeese, 1956), and well below the lethal limit for molting individuals and larvae (Scarrat & Raine, 1967; Cobb, 1976; Aiken & Waddy, 1986). Although heavy benthic invertebrate mortalities, including lobsters, have been reported in several estuaries after particularly heavy spring runoffs (Thomas, 1968; Thomas & White, 1969), including in the Great Bay Estuary (Nelson, pers. comm.), it is equally important to understand the survival strategies utilized in normal years. Previous investigations indicate that lobsters may use behavioral mechanisms to avoid low salinity conditions (Munro & Therriault, 1983; Reynolds & Casterlin, 1985; Vetrovs, 1990; Howell & Watson, 1991; Maynard, 1991; Watson & Howell, 1991). In this study, we measured the hemolymph osmolarity, oxygen consumption, heart rate and ventilation rate of lobsters under conditions similar to the Great Bay Estuary during a typical spring runoff, to determine the magnitude of the physiological stress these conditions evoke, and thus the extent to which the osmoregulatory abilities of lobsters might limit their range or influence their distribution.

The lethal levels of salinity, temperature, and oxygen tension for lobsters acclimated to various combinations of these factors have been previously determined (McLeese, 1956; Charmantier et al., 1988) and imply that some intermolt adults and juveniles may be able to tolerate the range of salinities normally found within the estuary. Under "optimal" acclimation conditions in the laboratory (5 °C, 30 ppt salinity, 6.4 mg/l oxygen) the lethal salinity may be as low as 6 ppt. However, this is an unusual case and the lethal salinity is, on average, greater than 11 ppt (McLeese, 1956). McLeese also found that as the temperature increases above 20 °C, tolerance for low salinity decreases up to as high as 16.4 ppt for animals acclimated to approximately 25 °C. Therefore, it would be expected that a low salinity event that occurred in the late spring/summer, when temperatures may reach as high as 25 °C, would be more stressful than one occurring in seasons when the water is cooler.

Most studies of juvenile and adult *H. americanus* metabolism, both in terms of oxygen consumption (McLeese, 1964; McMahon & Wilkens, 1975; Penkoff & Thurberg, 1982) and heart and ventilation rates (Wilkens & McMahon, 1972; Bill & Thurberg, 1985) have been conducted under conditions of invariable, "normal" salinity (\approx 30-33 ppt). Their metabolic rate rises with increasing activity, feeding, and temperature (McLeese,

ds are considered to be poor extends into many bays and te, 1969; Briggs & Mushacke, 1985; Estrella & McKiernan, 1991; Robichaud & Campbell, characterized by large seasonal m melting snow, which limits appropriate behavioral and/or , 1975; Vernberg & Vernberg,

ops to between 10 and 15 ppt 3), which is close to the lethal ow the lethal limit for molting (1976; Aiken & Waddy, 1986). g lobsters, have been reported ffs (Thomas, 1968; Thomas & on, pers. comm.), it is equally in normal years. Previous mechanisms to avoid low salinity asterlin, 1985; Vetrovs, 1990; owell, 1991). In this study, we tion, heart rate and ventilation Estuary during a typical spring stress these conditions evoke, lobsters might limit their

tension for lobsters acclimated eviously determined (McLeese, ermolt adults and juveniles may und within the estuary. Under 5 °C, 30 ppt salinity, 6.4 mg/l owever, this is an unusual case ppt (McLeese, 1956). McLeese 0 °C, tolerance for low salinity mated to approximately 25 °C. event that occurred in the late is 25 °C, would be more stressful er. abolism, both in terms of oxygen 1975; Penkoff & Thurberg, 1982) n, 1972; Bill & Thurberg, 1985) 'normal' salinity (\approx 30-33 ppt). ling, and temperature (McLeese,

1964; Leavitt, 1987) and also as molting approaches (McLeese, 1956; Penkoff & Thurberg, 1982). However, heart and ventilation rates in *H. americanus* are not affected by moderate reductions in ambient oxygen levels (McMahon & Wilkens, 1975). To our knowledge, no published data are presently available concerning the effect of low salinity on metabolic rate in *H. americanus*.

Several studies have shown that decreasing salinity effects the metabolic rate of crustaceans (Table 1). *Carcinus maenas* shows an increase in oxygen consumption as salinity is reduced (Taylor, 1977) presumably due to the increased energy necessary to actively pump ions. In the amphipod, *Onisimus glacialis*, increasing oxygen consumption was also correlated with decreasing salinity. The authors conclude that "It is questionable whether amphipods could endure such an increase in metabolic rate for long periods, and migration into seawater with higher salinity might be necessary for survival (Aarset & Aunaas, 1990)."

While the American lobster is a poor osmoregulator, in habitats with reduced salinity it does maintain its hemolymph osmolarity higher than the ambient environment. In this study our aim was to indirectly measure how much energy this limited osmoregulation requires, and then use this information to judge if the most energetically efficient adaptation to low salinity is increased active transport of ions or avoidance behavior. We found that lobsters consume more than twice as much oxygen at 10 ppt as compared to 20 ppt which would cause considerable stress during prolonged seasonal drops in salinity. We suggest that behavioral strategies would be more adaptive, under these circumstances, than extended periods of actively pumping ions. Furthermore, we discovered that at 10 ppt female lobsters require more energy than males to maintain the same hemolymph osmolarity. This would make the upper estuary even more stressful to females than it is to males, and may be one of the underlying causes for the observed

Table 1
Changes in metabolic rate for several decapod crustaceans exposed to low salinity

Oxygen consumption			Species	Source	Comments
Δ in ppt	Δ in ml O ₂ /g/h	% increase/10 ppt			
30-5	0.0093-0.0178	36.5	<i>Panopeus herbstii</i>	Dimock & Groves, 1975	Acclimated to 30 ppt, 10 °C
30-15	0.205-0.293	28.6	<i>Penaeus japonicus</i>	Chen & Lai, 1993*	Juveniles at 15 °C
37-10	0.20-0.40	37.0	<i>Penaeus japonicus</i>	Dalla Via, 1986	Approx. values
34-12	0.030-0.052	30.3	<i>Carcinus maenus</i>	Taylor, 1977	Approx. values
30-15	0.028-0.036	19.0	<i>Carcinus maenus</i>	Taylor et al., 1977	Tested at 14 °C
26-2	0.072-0.148	44.0	<i>Callinectes rathbunae</i>	Rosas et al., 1989	
35-5	1.6-3.4	37.5	<i>Callinectes sapidus</i>	Engel & Eggert, 1974	Approx. values from excised gill
30-10	0.138-0.175	13.4	<i>Callinectes sapidus</i>	Findley et al., 1978	Approx. values acclimated to 30 ppt, 20 °C

* See for review of salinity effects on oxygen consumption rates for penaeid shrimp.

domination of the upper Great Bay estuary by male lobsters (Vetrovs, 1990; Howell & Watson, 1991).

2. Methods

2.1. Animals

Animals were collected using standard vinyl-coated wire traps in the Great Bay Estuary, between March and August of 1991 and August and September of 1992. They were held at the Jackson Estuarine Laboratory, Durham, NH, which gets its water from the Great Bay Estuary, until 2 wk before testing, when they were transferred to recirculating aquaria (15 °C, 20 ppt) at the University of New Hampshire Zoology Department. Only adult (75–92 mm CL) intermolt lobsters, stages C4-D (Aiken, 1980), were used because molting appears to increase oxygen consumption and make animals less tolerant of low salinity (Penkoff & Thurberg, 1982). In addition, lobsters were not fed for at least 48 h prior to trials, and the activity of the animals was minimized during physiological testing (McLeese, 1964; Spoek, 1974).

Twenty ppt was used as a baseline salinity level for the following reasons: (1) previous work by McLeese (1956) demonstrated that 100% of the lobsters exposed to this salinity were able to survive; (2) our preliminary experiments indicated that when animals were dropped from 30 to 20 ppt, there was relatively little change in oxygen consumption; and (3) we were attempting to mimic the type of stress they would encounter during a typical spring in the estuary, when the salinity commonly drops from ≈ 20 to 10 ppt.

2.2. Hemolymph osmolarity experiments

Animals of both sexes were housed together (six per experiment) in a recirculating, refrigerated tank maintained at 15 °C and 20 ppt. They were allowed at least 72 h to acclimate to this environment (Dall, 1970). Within this tank two smaller tanks, one at 15 ppt and one at 10 ppt, were inserted so that the temperatures in all three tanks were identical. Salinities in the tanks were checked daily using an osmometer and a temperature compensated refractometer. The experiments were designed to mimic the salinity conditions in the respirometry experiments described below, and thus provide hemolymph osmolarity data that could be compared with changes in oxygen consumption at equivalent salinity values. The protocol consisted of moving animals between tanks so that they consecutively spent 24 h at 20, 15, and 10 ppt. After 24 h in 10 ppt, three animals, chosen randomly, were placed in the 15 ppt tank for 24 h and then in the 20 ppt tank. The other three animals remained in the 10 ppt tank. Hemolymph samples were taken sequentially after 1, 3, 6, and 24 h in each successive salinity to determine how the hemolymph osmolarity changes over time. In a similar experiment on *Panopeus herbstii* it was found that repeated sampling of the same individual compared to sampling each individual only once yielded similar results (Blasco & Forward, 1988). Approximately 1 ml of hemolymph was collected from the arthroidal membranes

at the base of the walking legs using a chilled syringe. Samples were immediately placed on ice to prevent clotting. Osmolarity was measured within 20 min with a Wescor 5100C Vapor Pressure Osmometer.

2.3. Oxygen consumption experiments

The oxygen consumption of single, unrestrained lobsters was measured in a continuous flow, open respirometer (Fig. 1). Movement was kept to a minimum by using a small (3.5" diameter) chamber, similar in size and shape to burrows found in the field (Atkinson & Taylor, 1988). The animal chamber was covered with clear, red plastic (Wald & Hubbard, 1957), and a black curtain was drawn in front of the entire apparatus to reduce visual disturbance of animals during experiments. However, the clear ends of the tube were left uncovered so that animals could be exposed to a normal photoperiod. A water jacket surrounded the 2-l animal chamber and the 100 ml oxygen electrode chambers to maintain their temperature at 15 °C. A temperature of 15 °C was chosen because it was equivalent to the average late spring temperature in the estuary (Loder et al., 1983).




Fig. 1. Design of the flow-through respirometer. Seawater of 20, 15, or 10 ppt flowed sequentially from the ambient seawater reservoir to the "in" electrode chamber, the animal chamber, the "out" electrode chamber, and finally back to the reservoir where it was continuously aerated. All connections between the chambers were made with three-way stopcocks so that the animal chamber could be bypassed to recalibrate or zero the oxygen electrodes. The heart, scaphognathite, and common ground electrodes were run through a rubber stopper in the endcap that was sealed with two O-rings and then placed into one end of the animal chamber. Cardiac and ventilation activity were recorded on a polygraph, while the outputs of the two oxygen electrodes were digitized, stored and displayed on a computer. Note that the electrode and animal chambers were jacketed as shown, however, the plumbing for the recirculating 15 °C seawater, used to maintain these chambers at a constant temperature, is not shown for the sake of simplicity.

Seawater with a salinity of 10, 15 or 20 ppt was pumped from a 15 °C holding tank to the electrode and animal chambers using a Masterflex peristaltic pump. The oxygen solubility values used for the different salinities were 20 ppt = 6.30 ml/l, 15 ppt = 6.50 ml/l, 10 ppt = 6.70 ml/l (Green & Carritt, 1967). The rate of flow varied between individuals, from ≈ 120 –160 ml/min (7.2–9.6 l/h), depending on the oxygen demand of the animal. The flow was set in the control run, and kept constant throughout the experiment, so that the oxygen concentration of the water leaving the animal chamber was no less than half the value of the water entering the chamber (≈ 3 ml/l). This prevented the chamber from becoming hypoxic, which would have had additional affects on the physiology of the animal (MacMahon & Wilkens, 1975). Transitions from one salinity to the next occurred at the rate of ≈ 0.20 ppt/min. Salinities of the outflow were monitored intermittently using a temperature compensated refractometer or a conductivity meter.

The oxygen tension of the water entering and leaving the chambers was measured with Strathkelvein polarographic oxygen electrodes (electrode model no. 1302; meter model no. 781). The analog outputs of the oxygen meters were digitized, stored, and displayed on a computer using an 8-channel analog-digital converter and the oscilloscope simulation program MacScope. The oxygen concentration was measured initially in the "in" electrode chamber receiving completely oxygenated water. The water then ran from this chamber to the animal chamber and the oxygen concentration of the water leaving the animal chamber was measured by the "out" electrode, located in the out electrode chamber. These values could then be subtracted to determine how much oxygen the lobster extracted from the water as it passed through the animal chamber. These measurements were then converted to oxygen consumption values using the following equation:

$$\text{mlO}_2/\text{g/h} = \frac{\{[\text{O}_2^{\text{in}}(\text{ml/l}) - \text{O}_2^{\text{out}}(\text{ml/l})] \times \text{flow rate}(\text{l/h})\}}{\text{wt of animal (g)}}$$

where O_2^{in} = oxygen tension measured at the "in" electrode, and O_2^{out} = oxygen tension recorded at the "out" electrode. Weight (wt) of animal was obtained after each trial by shaking and blotting the animal dry and then weighing it on a triple beam balance. Weight loss during a trial was assumed to be negligible. Oxygen consumption measurements were taken after 24 h at each salinity and small variations were minimized by averaging the data over at least 30 min for any one "data point" at a specific salinity for any individual (Hayes et al., 1992).

Following at least 2 wk of acclimation at 20 ppt, lobsters were placed in the animal chamber, and allowed to acclimate to the apparatus for at least another 24 h at 20 ppt and 15 °C. Sample sizes for subsequent analyses were as follows: (1) oxygen consumption of animals sequentially exposed to 20, 15, and 10 ppt ($n = 11$); (2) heart and scaphognathite rates of animals exposed to 20, 15, and 10 ppt ($n = 7$); (3) oxygen consumption, heart, and scaphognathite rates of animals spending one day at each progressive salinity from 20, to 15, to 10, back up to 15, and then returning to 20 ppt ($n = 4$). Lobsters remained in the respirometer for the entire experiment which lasted from 3–5 days. Most of the data illustrated in this paper was obtained during the final hour spent at each salinity; that is, after 24-h exposure to a given salinity.

2.4. Ventilation and heart rates

Oxygen consumption was used to indirectly measure relative changes in metabolic rate and throughout this manuscript we use the terms interchangeably. However, the two terms are not identical and it is certainly possible that there were times when our oxygen consumption measurements did not fully reflect 100% of the ongoing metabolic activity. Heart and ventilation rate are also sensitive measures of metabolic rate in crustaceans (Wilkins & MacMahon, 1972; MacMahon & Wilkins, 1975; Hume & Berlind, 1976; Cumberlidge & Uglow, 1977; Dyer & Uglow, 1977; Mercado-Allen & Thurberg, 1987; Hamilton & Houlihan, 1992). To confirm that changes in oxygen consumption were related to variations in oxygen uptake and circulation, we also monitored the heart and ventilation rates of several animals while they were in the respiration chambers. A pair of copper wire electrodes was implanted dorsal to the cervical groove to record heart activity and another pair was placed laterally above one of the branchial chambers to monitor scaphognathite movements. Signals were amplified and displayed on a Grass Model 7D Polygraph, and simultaneously transformed into instantaneous rate by a tachograph. The output of the tachograph was displayed on the polygraph along with the original heart and scaphognathite signals, and also digitized and stored along with the oxygen consumption data.

To confirm that the signal recorded from the lateral electrodes was an accurate indicator of gill bailer activity, simultaneous recordings were obtained with the Grass Polygraph and a UFI model 2991 impedance transducer (Dyer & Uglow, 1977; Bill & Thurberg, 1985) using electrodes implanted at the same location in the carapace. This technique verified that the signal was representative of water flow through the branchial chamber and thus a valid indicator of ventilation.

Heart and scaphognathite activity, as well as oxygen consumption, were measured after 24 h at each successive salinity. Measurements were also taken for 1 h prior to, and 2 h after, the transition to the next salinity. To assess any short term effects of salinity change during the transition period, short samples of heart and scaphognathite rate were taken throughout each trial.

3. Results

In both male and female lobsters, decreases in hemolymph osmolarity paralleled reductions in ambient osmolarity (Fig. 2). There were no significant differences between the sexes in their ability to maintain their hemolymph slightly hyperosmotic to the surrounding seawater ($p > 0.70$, independent t -tests). However, the lobsters always regulated their internal osmolarity at a level that was 50–100 mOsm higher than ambient levels ($p < 0.05$ for both sexes at all salinities tested, except for females at 20 ppt, where $p = 0.075$, independent t -tests). Thus, as shown by Dall (1970), lobsters are limited osmoregulators at salinities below 20 ppt.

The degree to which lobsters osmoregulate seems to be related to the amount of osmotic stress they encounter. At salinities of 20–30 ppt there is comparatively little difference between their internal osmolarity and the osmolarity of the surrounding




Fig. 2. Hemolymph osmolarity as a function of ambient osmolarity. Samples were obtained after 24-h exposure to each osmolarity in succession, from 20 to 15 to 10 ppt. Below 20 ppt, lobsters maintain the internal osmolarity at levels above ambient salinity ($p < 0.05$ for all animals except females at 20 ppt where $p = 0.075$, independent t -tests), but the hemolymph concentration continues to decrease as the salinity drops. Thus, they are considered limited osmoregulators. Each point represents the mean of either 12 males or 16 females \pm SEM.

seawater (Dall, 1970). However, as the salinity is reduced toward 10 ppt, the difference between the ambient and hemolymph osmolarities increases. As the salinity drops, most of the concurrent drop in hemolymph osmolarity occurs within the first hour and then stabilizes at a lower, albeit hyperosmotic, level after ≈ 6 h (Fig. 3), which is similar to the value of 4 h found for *P. herbstii* exposed to an 8 ppt drop in ambient salinity (Blasco & Forward, 1988). Lobsters are able to maintain this osmolarity set point for several days, but as shown below, this requires a considerable expenditure of energy at low salinities.

There was a close relationship between oxygen consumption, scaphognathite beat and heart rate (Fig. 4). This has also been demonstrated in a number of other marine




Fig. 3. Changes in hemolymph osmolarity over time. Hemolymph samples were taken after 1, 3, 6 and 24 h at each successive salinity, from 20 to 10 ppt. The most significant drop occurs within the first hour of exposure to the next lower salinity, and there is little change between 6- and 24-h exposure. Each point represents the mean \pm SEM ($n = 23$); except at 25 h ($n = 6$) and 54 h ($n = 12$).




Fig. 4. A continuous 2-h record of oxygen consumption, heart rate, and ventilation rate, recorded from an adult lobster, at 20 ppt and 15 °C. There is a close relationship between oxygen consumption, heart and scaphognathite rates. Heart and scaphognathite activity, along with pO₂ values from the respirometer, were obtained as described in the Methods section. In order to induce large fluctuations in respiration and circulation the outside of the respirometer was intermittently tapped. Note that there is a slight delay between changes in ventilation and heart rates and changes in oxygen consumption. This is due to the delay inherent in the flow-through respirometer.

crustaceans (*Cancer pagurus*: Ansell, 1973; *H. gammarus*: Spoek, 1974; *C. maenus*: Cumberland & Uglow, 1977; Taylor, 1977; Hamilton & Houlihan, 1992). Covariation of cardiac and ventilatory rhythms in *H. americanus* has been reported previously (MacMahon & Wilkens, 1975; Mercaldo-Allen & Thurberg, 1987) and was observed in these experiments as well (Fig. 4). For example, during periods of little or no scaphognathite activity and associated bradycardia, there were clear drops in oxygen consumption, whereas oxygen consumption increased when heart and ventilation rates were elevated. In *C. maenus* there is a closer correlation between oxygen consumption and ventilation than oxygen consumption and heart rate (Cumberland & Uglow, 1977; Taylor, 1977; Hamilton & Houlihan, 1992) and this appears to be true for *H. americanus* as well (Fig. 5). Thus, for certain types of studies, once the precise relationship has been established, heart or ventilation rates could be used to indirectly monitor oxygen consumption.

Exposure of lobsters to low salinities, representative of those found in the estuary, caused an increase in their oxygen consumption, heart, and scaphognathite rates. As the ambient salinity was reduced from 20 to 15 to 10 ppt there was a nearly linear increase in oxygen consumption (Fig. 6), with significant differences between the metabolic rates at each salinity ($n = 11$) ($p < 0.001$, repeated measures ANOVA; Scheffe's F -test). There were also significant differences in heart and scaphognathite rates between 20 and 10 ppt ($n = 7$) ($p < 0.01$, repeated measures ANOVA; Scheffe's F -test) but these differences were not significant between 20 and 15, or 15 and 10 ppt. As the salinity was gradually restored to 20 ppt, oxygen consumption, heart and scaphognathite rates approached the levels found prior to the "low salinity event" (Fig. 7 A,B).

Although male and female lobsters showed a similar ability to maintain their




Fig. 5. Correlation of lobster oxygen consumption, heart and scaphognathite rates. Both heart ($Y = 24.04 + 658.24X$, $R^2 = 0.37$) and ventilation ($Y = 17.21 + 1969.20X$, $R^2 = 0.60$) rates were associated with changes in oxygen consumption. All values were taken from lobsters which had been at a particular salinity (20, 15, or 10 ppt), for 24 h. Data points represent values which were averaged over ≈ 30 -min intervals. Data from seven different lobsters were used for this plot to provide a wide range of values.

hemolymph slightly hyperosmotic to the surrounding seawater (Fig. 2), at low salinities there exists an apparent sex related difference in the amount of energy required for this osmoregulatory activity (Fig. 8). There is no significant difference between male ($n = 6$) and female ($n = 5$) percent increase in oxygen consumption at 15 ppt, but at 10 ppt there is a strong trend for the proportional increase in metabolic rate of female: to be greater than that observed for males (presumably due to differential osmoregulatory abilities) ($p = 0.08$, independent t -test on log-transformed data). The slopes of the regression lines for the percent increase in metabolic rate at 15 and 10 ppt, from 20 ppt were also significantly different between males and females (ANCOVA, $p < 0.01$). Upon recovery, males ($n = 2$) consumed 0.039 ± 0.0118 ml $O_2/g/h$ at 15 ppt and 0.026 ± 0.0041 ml $O_2/g/h$ at 20 ppt, whereas females ($n = 2$) consumed 0.057 ± 0.003




Fig. 6. The relationship between oxygen consumption and ambient salinity. Salinity was gradually dropped in 5 ppt increments each day, from 20 to 15 to 10 ppt. Test animals ($n = 11$) had significantly greater oxygen consumption at each decreasing salinity ($p < 0.001$, repeated measures ANOVA; Scheffe's F -test).




Fig. 7. Oxygen consumption (A), as well as heart and ventilatory (B), responses during decreases, and subsequent increases, in ambient salinity levels. The oxygen consumption, heart rate and scaphognathite rates (\pm SEM) of mature lobsters ($n=4$) were measured after 24 h of continuous exposure to the following sequence of salinities: 20, 15, 10, 15, and 20 ppt. It is likely that the changes in oxygen consumption, heart rate, and ventilation rate recorded from these otherwise quiescent lobsters are a manifestation of changing osmoregulatory demands.

and 0.045 ± 0.0022 at the same salinities. Because of small sample sizes the significance of this data is unclear, but it seems to indicate that females recovered more slowly than the males as the salinity was subsequently increased. Thus, at salinities below 15 ppt it appears as though male lobsters are more efficient osmoregulators than female lobsters.

4. Discussion

Lobsters and other estuarine invertebrates encounter large seasonal drops in salinity that may last from several days to weeks. Previous studies (Dall, 1970), and those reported here, indicate that lobsters are only limited osmoregulators, allowing their hemolymph osmolarity to drop, but always maintaining it 50-100 mOsm higher than their ambient environment. Osmoregulation in lobsters appears to require a significant amount of energy, especially at the low salinities characteristic of estuarine habitats in a typical spring. In our experiments, at 20 ppt, lobsters consumed 0.029 ± 0.003 ml of oxygen/g/h, which is very similar to the values reported in the literature at compa-

both heart associated particular \approx 30-min values.

w salini- up for be ale t, but at f females smoregu- es of the n 20 ppt, > 0.01). ppt and ± 0.0037

dropped, water oxy-